

# An H3K36 Methylation-Engaging Tudor Motif of Polycomb-like Proteins Mediates PRC2 Complex Targeting

Ling Cai,<sup>1,2,9</sup> Scott B. Rothbart,<sup>1,2,9</sup> Rui Lu,<sup>1,2,9</sup> Bowen Xu,<sup>1,2,3</sup> Wei-Yi Chen,<sup>4</sup> Ashutosh Tripathy,<sup>2</sup> Shira Rockowitz,<sup>6</sup> Deyou Zheng,<sup>6</sup> Dinshaw J. Patel,<sup>7</sup> C. David Allis,<sup>5</sup> Brian D. Strahl,<sup>1,2,3</sup> Jikui Song,<sup>8,\*</sup> and Gang Greg Wang<sup>1,2,3,\*</sup>

<sup>1</sup>UNC Lineberger Comprehensive Cancer Center

<sup>2</sup>Department of Biochemistry and Biophysics

<sup>3</sup>Biological and Biomedical Sciences Program

University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC 27599, USA

<sup>4</sup>Laboratory of Biochemistry and Molecular Biology

<sup>5</sup>Laboratory of Chromatin Biology and Epigenetics

The Rockefeller University, New York, NY 10065, USA

<sup>6</sup>Department of Neurology, Albert Einstein College of Medicine, Bronx, NY 10461, USA

<sup>7</sup>Structural Biology Program, Memorial Sloan-Kettering Cancer Center, New York, NY 10065, USA

<sup>8</sup>Department of Biochemistry, University of California, Riverside, CA 92521, USA

<sup>9</sup>These authors contributed equally to this work

\*Correspondence: [jikui.song@ucr.edu](mailto:jikui.song@ucr.edu) (J.S.), [greg\\_wang@med.unc.edu](mailto:greg_wang@med.unc.edu) (G.G.W.)

<http://dx.doi.org/10.1016/j.molcel.2012.11.026>

## SUMMARY

Polycomb repressive complex 2 (PRC2) regulates pluripotency, differentiation, and tumorigenesis through catalysis of histone H3 lysine 27 trimethylation (H3K27me<sub>3</sub>) on chromatin. However, the mechanisms that underlie PRC2 recruitment and spreading on chromatin remain unclear. Here we report that histone H3 lysine 36 trimethylation (H3K36me<sub>3</sub>) binding activity is harbored in the Tudor motifs of PRC2-associated polycomb-like (PCL) proteins PHF1/PCL1 and PHF19/PCL3. Ectopically expressed PHF1 induced Tudor-dependent stabilization of PRC2 complexes on bulk chromatin and mediated spreading of PRC2 and H3K27me<sub>3</sub> into H3K36me<sub>3</sub>-containing chromatin regions. In murine pluripotent stem cells, we identified coexistence of H3K36me<sub>3</sub>, H3K27me<sub>3</sub>, and PHF19/PCL3 at a subset of poised developmental genes and demonstrated that PHF19/PCL3 Tudor function is required for optimal H3K27me<sub>3</sub> and repression of these loci. Collectively, our data suggest that PCL recognition of H3K36me<sub>3</sub> promotes intrusion of PRC2 complexes into active chromatin regions to promote gene silencing and modulate the chromatin landscape during development.

## INTRODUCTION

Modulation of the chromatin landscape by covalent histone posttranslational modifications (PTMs) represents a fundamental way of regulating DNA-templated processes such as gene

transcription (Chi et al., 2010; Kouzarides, 2007). Histone H3 lysine 27 trimethylation (H3K27me<sub>3</sub>) promotes gene silencing, whereas promoter-associated histone H3 lysine 4 trimethylation (H3K4me<sub>3</sub>), together with gene body-associated H3 lysine 36 trimethylation (H3K36me<sub>3</sub>), demarcates active genes (Mikkelsen et al., 2007). PRC2 complexes are the major enzymatic machineries responsible for writing H3K27me<sub>3</sub>, and PRC2-mediated gene silencing is involved in various biological processes including stem cell pluripotency, differentiation, and cancer progression (Bracken and Helin, 2009; Margueron and Reinberg, 2011). While a reconstituted tetrameric PRC2 core complex (comprising EZH2, EED, SUZ12, and NURF55/RbAp48) is sufficient to induce H3K27me<sub>3</sub> (Cao and Zhang, 2004; Margueron and Reinberg, 2011), a number of PRC2-associated factors, including EED (Margueron et al., 2009), JARID2 (Li et al., 2010; Pasini et al., 2010; Peng et al., 2009; Shen et al., 2009), and noncoding RNAs (Bracken and Helin, 2009; Margueron and Reinberg, 2011), that either modulate H3K27me<sub>3</sub> catalysis or help to stabilize and tether PRC2 to appropriate genomic regions have recently been identified. Despite these recent advances, mechanisms responsible for PRC2 targeting and spreading on active chromatin to promote de novo repressed states remain poorly defined.

In this study, we show that two PCL family proteins (PHF1/PCL1 and PHF19/PCL3), accessory components of the PRC2 core complex (Cao et al., 2008; Hunkapiller et al., 2012; Sarma et al., 2008), harbor H3K36me<sub>3</sub>-reading activity within their N-terminal Tudor motifs. Biochemical, biophysical, and structural analyses reveal tight binding to H3K36me<sub>3</sub> through a conserved aromatic cage formed by PCL Tudor domains. Our gene regulation and genomics analysis, using both over-expression and knockdown systems, define the PCL Tudor-H3K36me<sub>3</sub> interaction as critical for both the targeting and spreading of PRC2 into active chromatin regions and for the maintenance of optimal repression of poised developmental

genes where PCL, H3K36me<sub>3</sub>, and H3K27me<sub>3</sub> coexist. Our studies shed important light on the regulation of PRC2 by PCL proteins, implicating H3K36me<sub>3</sub> reading in this process, and contribute to our understanding of the dynamics of gene expression and chromatin remodeling associated with development and disease (Chi et al., 2010).

## RESULTS

### PHF1 Directly Binds to H3K36me<sub>3</sub> through a Conserved N-Terminal Tudor Motif

We began this study by attempting to identify, by mass spectrometry, H3K36me<sub>3</sub>-reader proteins using synthetic histone tail peptides as bait for nuclear extracts. In addition to the previously identified putative H3K36me<sub>3</sub>-reading effectors NSD2 and MSH6 (Vermeulen et al., 2010), we identified PHF1 (Figures 1A and S1A), a known accessory component of H3K27me<sub>3</sub>-promoting PRC2 complexes (Cao et al., 2008; Sarma et al., 2008). PHF1 and its homologs MTF2/PCL2 and PHF19/PCL3 have several putative chromatin-interacting domains (Cao et al., 2008) including a highly conserved N-terminal Tudor domain and two plant homeodomain (PHD) fingers (Figure S1B). In-solution peptide pull-down assays with recombinant GST-fusion proteins of each of these individual domains of PHF1 indicated that the Tudor domain facilitated H3K36me<sub>3</sub> recognition (Figures 1B and S1C). To confirm this interaction, we established stable cell lines expressing wild-type (WT) or N-terminal truncated forms of PHF1 lacking either the Tudor domain or PHD fingers. Indeed, WT PHF1, but not that lacking the Tudor domain, pulled down with H3K36me<sub>3</sub> peptides (Figures 1C and S1D). Furthermore, PHF1 coimmunoprecipitated (coIP) with endogenous histones that contained H3K36me<sub>3</sub> in a Tudor-dependent manner (Figure 1D). Collectively, our data implicated the PHF1 Tudor domain (PHF1<sub>Tudor</sub>) as an H3K36me<sub>3</sub>-reading effector module.

### PHF1<sub>Tudor</sub> Binds Specifically and with High Affinity to H3K36me<sub>3</sub>


We next sought to determine the histone-binding specificity of PHF1<sub>Tudor</sub> in an unbiased manner by using a recently developed high-density histone peptide microarray platform (Fuchs et al., 2011; Rothbart et al., 2012b). Arrays were printed with a library of unmodified and modified peptides (Table S1) containing known single and combinatorial histone PTMs (methylation, acetylation, and phosphorylation) on all four core histones (H2A, H2B, H3, and H4). Interrogation of the GST-tagged PHF1<sub>Tudor</sub> by peptide arrays (Figure 1E, top panel) revealed a strong preference for various H3K36me<sub>3</sub>-containing peptides (Figures 1E and S1E). In addition, PHF1<sub>Tudor</sub> displayed weak interactions with other methylated histone peptides such as H3 lysine 9 trimethylation (H3K9me<sub>3</sub>) and H3K27me<sub>3</sub> (Figures 1F, S1F and S1G). Similar results were obtained using another peptide array platform (Figures S1H and S1I). Using isothermal titration calorimetry (ITC), we next quantified these observed interactions. Consistent with array results and peptide pull-downs, ITC measurements of the PHF1<sub>Tudor</sub> demonstrated high affinity and specificity for H3K36me<sub>3</sub>, with a dissociation constant ( $K_D$ ) of  $\sim 4.2$   $\mu$ M,  $\sim 19$   $\mu$ M, and  $\sim 38$   $\mu$ M for tri-, di-,

and monomethylated H3K36 states, respectively (Figure 1G, Table 1, and Figures S1L–S1O). Compared to H3K36me<sub>3</sub>, interactions with H3K9me<sub>3</sub> and H3K27me<sub>3</sub> were significantly weaker, with a  $K_D$  of  $\sim 38$   $\mu$ M and  $\sim 168$   $\mu$ M, respectively (Figures S1P and S1Q and Table 1). We further determined the histone-binding specificity toward nucleosomal substrates in a GST pull-down assay using GST-PHF1<sub>Tudor</sub> and mononucleosomes purified from 293 cells. Again, PHF1<sub>Tudor</sub> preferentially bound to nucleosomes containing H3K36me<sub>3</sub>—but not other examined methylations, including those marked with H3K27me<sub>3</sub>, H3K9me<sub>3</sub>, H3K4me<sub>3</sub>, trimethylation of H4 lysine 20, or asymmetric dimethylation of H3 arginine 2 and arginine 17 (Figure 1H, lane of WT). Collectively, these data showed that PHF1<sub>Tudor</sub> motif demonstrates tight affinity and high selectivity toward H3K36me<sub>3</sub>.

Full-length PHF1 localizes exclusively in the nucleus (Figure 3D). We therefore asked whether association of full-length PHF1 with chromatin is dependent on H3K36me<sub>3</sub>. To do this, we diminished global levels of H3K36me<sub>3</sub>/H3K36me<sub>2</sub> on chromatin by overexpressing an H3K36me<sub>3</sub>/H3K36me<sub>2</sub> demethylase JMJD2A/KDM4A in 293 cells stably expressing a FLAG-tagged form of PHF1. Consistent with previous findings (Klose and Zhang, 2007), we observed a dramatic reduction in H3K36me<sub>3</sub>/H3K36me<sub>2</sub> and H3K9me<sub>3</sub> after biochemical separation of JMJD2A/KDM4A-overexpressing cells into chromatin-bound and soluble cytoplasmic/nucleoplasm fractions with no change detected in H3K4me<sub>3</sub> or H3K27me<sub>3</sub> (Figure 1I). Importantly, compared to control, a reduction was observed in the chromatin-bound fraction of PHF1, concurrent with an increase in its soluble pool (Figure 1I), thus demonstrating that PHF1 is less stable on chromatin in the absence of global H3K36me<sub>3</sub>/H3K36me<sub>2</sub>. Collectively, our in vitro and cellular biochemical data comprehensively showed that PHF1<sub>Tudor</sub> specifically and strongly binds to H3K36me<sub>3</sub> and that this interaction is necessary to target PHF1 to chromatin. In addition, the PHF1<sub>Tudor</sub>-H3K36me<sub>3</sub> interaction represents the first high-affinity effector interaction to H3K36me<sub>3</sub>, which is several orders of magnitude tighter than all of the previously reported H3K36me<sub>3</sub>-reading motifs (Sun et al., 2008; Vezzoli et al., 2010; Xu et al., 2008).

### Structural Analysis of PHF1<sub>Tudor</sub> Revealed a Unique H3K36me<sub>3</sub>-Reading Pocket

To gain insight into the molecular basis for recognition of H3K36me<sub>3</sub> by PHF1<sub>Tudor</sub>, we performed NMR spectroscopy and obtained the solution structure of PHF1<sub>Tudor</sub> (residues 7–83) in complex with an H<sub>3</sub><sub>31–41</sub>K36me<sub>3</sub> peptide. Initial comparison of <sup>1</sup>H, <sup>15</sup>N-HSQC spectra of the unbound PHF1<sub>Tudor</sub> with those in the presence of H3K36me<sub>3</sub> indicated that, while the majority of PHF1<sub>Tudor</sub> residues exhibited no detectable change in chemical shifts, several residues showed a large perturbation, indicating a role for these residues in histone tail engagement (Figure S2A). The NMR structures of the PHF1<sub>Tudor</sub>-H3K36me<sub>3</sub> complex revealed that residues 31–81 of PHF1<sub>Tudor</sub> adopted a classic Tudor domain fold (Figure 2A and Table 2) as initially observed in the Survival of Neuron Tudor domain (Selenko et al., 2001), whereas flanking residues were disordered, evidenced by poor chemical shift dispersion and lack of NOE connectivity. We also found that the structure of PHF1<sub>Tudor</sub> is



**Figure 1. The N-Terminal Tudor Motif of PHF1 Specifically Recognizes H3K36me3**

(A) Enrichment of PHF1 in pull-downs using biotinylated histone tail peptides that contain H3K36me3 as compared to that using unmodified H3K36. See also Figure S1.

(B) Pull-down of GST recombinant proteins fused to different PHF1 domains using H3K36me3 or unmodified H3K36 peptides (top two panels). Input is shown in bottom panels.

(C) Immunoblot following H3K36me3 peptide pull-down (top panel) from extracts of cells transduced with vector or that encoding the full-length PHF1 or deletion forms. Input of peptides and PHF1 is shown in middle and bottom panels.

(D) colP using FLAG antibodies detecting a Tudor-dependent association of PHF1 (middle) to H3K36me3-containing histones (top) in 293 cells.

(E) Representative scan images of histone peptide microarrays probed with either WT or mutant (Y47A) GST-PHF1 Tudor proteins. Highlighted are positions of H3K36me3-containing peptides (blue) and IgG control (yellow). See also Table S1.

(F) Scatter plot of two peptide arrays probed with the PHF1 Tudor motif. Peptides are colored according to the legend. All other peptides are shown in black. Correlation coefficient was derived from linear regression analysis using GraphPad Prism.

(G) ITC measurements of binding affinities of PHF1 Tudor to H3 peptides containing different H3K36 methylation states. Insert represents silver staining of 1% of input (lane 1) and pull-down samples of PHF1 Tudor (top panel) using peptides harboring un-, mono-, di-, or trimethylated H3K36 (lanes 2–5, bottom panel).

(H) Immunoblot using various histone methylation antibodies on mononucleosomes pulled down by either WT or mutant (Y47A) GST-PHF1 Tudor proteins.

(I) Immunoblot of chromatin-bound and soluble cell fractions following 48 hr transient expression of JMJD2A/KDM4A. Mock indicates vector control.

**Table 1. Summary of Thermodynamic and Curve-Fitting Parameters for ITC Assays Using Recombinant PHF1 or PHF19 Tudor Domain Proteins and Various Histone Peptides**

Protein	Peptide	$\Delta H$ (cal/mol)	$\Delta S$ (cal/mol/deg)	$K_a$ (M <sup>-1</sup> )	N	Chi2/DoF	$K_D$ ( $\mu$ M)
PHF1 <sup>Tudor</sup> , WT	H3 <sub>(28-43)</sub> K36me3	-5,626 ± 108	4.45	$2.38 \times 10^5 \pm 1.91 \times 10^4$	0.79 ± 0.01	5,415	4.2 ± 0.3
PHF1 <sup>Tudor</sup> , WT	H3 <sub>(28-43)</sub> K36me2	-3,994 ± 261	7.26	$5.18 \times 10^4 \pm 6.79 \times 10^3$	0.94 ± 0.04	4,338	19.3 ± 2.5
PHF1 <sup>Tudor</sup> , WT	H3 <sub>(28-43)</sub> K36me1	-817 ± 342	17.3	$2.63 \times 10^4 \pm 1.12 \times 10^4$	0.70 ± 0.23	658.3	38.0 ± 16.1
PHF1 <sup>Tudor</sup> , WT	H3 <sub>(28-43)</sub> K36me0	-	-	-	-	-	N.D.
PHF1 <sup>Tudor</sup> , WT	H3 <sub>(19-35)</sub> K27me3	-5,141 ± 469	-1.21	$5.93 \times 10^3 \pm 486$	1.03 ± 0.07	347.7	168 ± 13.8
PHF1 <sup>Tudor</sup> , WT	H3 <sub>(1-15)</sub> K9me3	-817 ± 343	17.3	$2.63 \times 10^4 \pm 1.12 \times 10^4$	0.70 ± 0.23	658.3	38.0 ± 16.2
PHF1 <sup>Tudor</sup> , W41A	H3 <sub>(28-43)</sub> K36me3	-1,355 ± 457	15.8	$3.32 \times 10^4 \pm 1.20 \times 10^4$	0.68 ± 0.18	1,456	30.1 ± 10.9
PHF1 <sup>Tudor</sup> , Y47A	H3 <sub>(28-43)</sub> K36me3	-	-	-	-	-	N.D.
PHF1 <sup>Tudor</sup> , F65A	H3 <sub>(28-43)</sub> K36me3	-	-	-	-	-	N.D.
PHF1 <sup>Tudor</sup> , F71A	H3 <sub>(28-43)</sub> K36me3	-	-	-	-	-	N.D.
PHF1 <sup>Tudor</sup> , E66K	H3 <sub>(28-43)</sub> K36me3	-6,471 ± 229	-1.94	$4.39 \times 10^4 \pm 3.76 \times 10^3$	0.60 ± 0.02	5,055	22.8 ± 2.0
PHF19 <sup>Tudor</sup> , WT	H3 <sub>(27-46)</sub> K36me3	-6,610 ± 267	0.466	$1.60 \times 10^5 \pm 1.65 \times 10^4$	0.58 ± 0.02	7,436	6.2 ± 0.6
PHF19 <sup>Tudor</sup> , WT	H3 <sub>(27-46)</sub> K36me1	-	-	-	-	-	N.D.
PHF19 <sup>Tudor</sup> , Y56A	H3 <sub>(27-46)</sub> K36me3	-	-	-	-	-	N.D.

N.D., not determined. See also Figures S1L–S1S and S2G.


dominated by a  $\beta$ -barrel formed by five antiparallel  $\beta$  strands and that association of PHF1<sup>Tudor</sup> with H3K36me3 is mediated by direct contacts to both H3K36me3 and several surrounding histone tail residues (Figures 2B and 2C). The trimethylammonium side chain of H3K36me3 fits into an aromatic cage at one end of the  $\beta$ -barrel formed by residues Trp41, Tyr47, Phe65, Ser69, Asp67, and Phe71 (Figure 2D). Notably, Trp41 and Phe71 are not conserved in the Tudor domain of *Drosophila* polycomb-like (Pcl) protein (Figure S2B), which was reported to lack the ability to bind to histone methylation (Friberg et al., 2010), suggesting the role of these aromatic residues in formation of a closed cage composition. The overall structure of the H3K36me3-binding cage found in PHF1<sup>Tudor</sup> is reminiscent of those previously reported for the BRPF1 PWWP domain (Vezzoli et al., 2010) and the Eaf3 chromodomain (for comparison, see Figures S2C–S2E) (Sun et al., 2008; Xu et al., 2008). Several distinct features of PHF1<sup>Tudor</sup> were also revealed by our analysis, mainly occurring at contacts to histone residues surrounding H3K36me3. First, a salt bridge formed between the H3K37 side chain and the carboxylate of E66 of PHF1<sup>Tudor</sup> (Figure 2C, a dashed line). Second, the H3P38 side chain interacted with a hydrophobic patch generated by L38 and L46 of PHF1<sup>Tudor</sup> (Figures 2B and 2C). Third, hydrophobic contacts were observed between the H3H39 side chain and L38 and L48 of PHF1<sup>Tudor</sup> (Figures 2B and 2C).

To examine the functional relevance of observed contacts to H3K36me3 recognition, we introduced individual alanine substitutions to each of the H3K36me3-caging residues. We found that these single point mutations abrogated the binding of PHF1<sup>Tudor</sup> to H3K36me3 peptides on arrays (Figure 1E, Y47A, bottom panel) and by ITC (Figure 2E, Table 1, and Figure S1R) and peptide pull-downs (Figures 2F and S2F, left panel). Point mutations also abrogated binding of full-length PHF1 to H3K36me3 peptides (Figure 2G) and to H3K36me3-containing histones as assayed by GST-pull-downs (Figure 1H; Y47A) and colPs (Figure 2H). Furthermore, disruption of the salt bridge to H3K37 by

mutating E66 of PHF1<sup>Tudor</sup> to K66 also caused a significant reduction in H3K36me3 binding ( $K_D = \sim 22.8 \mu$ M; Figures S1S and S2F and Table 1). Collectively, our structural and mutational analysis demonstrated that unlike other known H3K36me3 effector proteins, numerous van der Waals contacts and electrostatic interactions contribute to a high-affinity association of PHF1<sup>Tudor</sup> with H3K36me3.

#### PHF1<sup>Tudor</sup> Alters Localization of PHF1-PRC2 Complexes, but Not PRC2 Core Structure

PHF1 was reported to interact with PRC2 core complexes where PHF1 modulates the enzymatic activity of associated PRC2 core complexes in vitro (Cao et al., 2008; Sarma et al., 2008). To determine whether PHF1 primarily associates with PRC2, we used mass spectrometry analysis to identify PHF1-associated factors from 293 stable cell lines. The top unique hits of identified proteins were the four subunits of tetrameric PRC2 core complexes (SUZ12, EED, NURF55/RbAp48, and EZH2 or EZH1) (Figure 3A and Table S2), consistent with comigration of PHF1 and PRC2 core components observed in gel filtration chromatography analyses (Cao et al., 2008; Sarma et al., 2008). We next sought to determine whether the PHF1<sup>Tudor</sup> motif was needed for PRC2 complex formation. colPs with EZH2 and SUZ12 in cells stably expressing WT or Tudor-mutated forms of PHF1 showed that membership in the PRC2-PHF1 complex was not dependent on PHF1<sup>Tudor</sup> (Figures 3B, S3A, and S3B). Notably, while JARID2 was previously identified as associating with the PRC2 core complex (Li et al., 2010; Pasini et al., 2010; Peng et al., 2009; Shen et al., 2009), we did not detect JARID2 in either mass spectrometry analysis (Table S2) or colP experiments with PHF1 (Figure S3C). These results are consistent with previous proteomics analysis of JARID2-associated complexes that failed to identify PCL proteins such as PHF1 or MTF2 (Shen et al., 2009). These studies collectively indicate that PHF1 primarily assembles in protein complexes with PRC2, potentially to modulate PRC2 targeting via its



**Figure 2. Structural and Biophysical Analysis Revealed Intermolecular Interaction between PHF1<sup>TUDOR</sup> and H3K36me3-Containing Histones**

(A) Backbone superposition of 20 energy-minimized conformers representing the NMR structure of PHF1 Tudor in complex with an H3K36me3 peptide. For clarity, residues 1–30 and 82–83 of PHF1 Tudor and 31–34 of H3K36me3 peptide are not shown due to disorder. The PHF1 Tudor (cyan) and bound H3K36me3 peptide are shown in ribbon representation with side chains of H3K36me3-binding cage (red) and H3K36me3 (yellow) in stick representation. See also Figure S2. (B) Surface electrostatic representation of PHF1 Tudor bound to H3K36me3 peptide. (C) Representative structure of PHF1 Tudor-H3K36me3 complex with side chains involved in intermolecular contacts. The salt bridge formed between PHF1 Tudor E66 and H3K37 is depicted as a red dashed line. (D) Positioning of the H3K36me3 side chain within an aromatic cage of the indicated residues (pink) on the surface of PHF1 Tudor. (E) ITC measurements of binding affinities of WT or mutant PHF1 Tudor to H3K36me3 peptides. (F) Silver staining of pull-down samples (top panel) of WT or mutant (Y47A and F65A) forms of PHF1 Tudor using histone tail peptides that harbor either un-, mono-, di-, or trimethylated H3K36. Inputs of protein and peptide used are shown in middle and bottom panels. (G) Anti-FLAG immunoblot following peptide pull-down (top panel) from extracts of cells transfected with either empty vector or vector encoding a WT or mutant form of FLAG-tagged PHF1. (H) Western blots examining association of PHF1 (top) to endogenous H3K36me3-containing histones (middle) in HeLa cells that expressed a WT or mutant form of FLAG-tagged PHF1.

histone-reading ability, and that PCL proteins and JARID2 reside in distinct PRC2 subcomplexes.

We next sought to determine whether the PHF1<sup>Tudor</sup> domain was required for PRC2 recruitment to chromatin. To this end, we fractionated PHF1 stable expression cells into chromatin and soluble fractions and found that, consistent with our observation following JMJD2A/KDM4A overexpression (Figure 1I), chromatin association of PHF1 was perturbed in the absence of Tudor recognition of H3K36 methylation (Figure 3C, lane 1 versus lanes 2 and 3). Notably, PRC2 association with bulk chromatin was also hindered in the absence of a functional PHF1<sup>Tudor</sup> in our PHF1 overexpression system, evidenced by a loss of SUZ12 and EZH2 from the chromatin fraction and a concurrent increased soluble pool of these PRC2-associated factors (Figure 3C). Consistent with these results, confocal

immunofluorescence microscopy of WT PHF1 showed punctate nuclear foci overlapping with DAPI-dense regions of chromatin, while Tudor mutant forms of PHF1 showed diffuse distribution throughout the nucleus with accumulations in the chromatin-free nucleoplasm (Figures 3D and S3D). Collectively, our data suggest that PHF1 alters chromatin localization or stabilization of PHF1 and associated PRC2 proteins via an H3K36me3/H3K36me2-recognizing PHF1<sup>Tudor</sup>, a function separated from assembly of a higher-order PHF1-PRC2 complex.

#### PHF1 Induces Spreading of PRC2 and H3K27me3 into H3K36me3-Marked Loci

Our data presented thus far suggest that PCL family proteins may link PRC2 core complexes to active chromatin via H3K36me3 recognition and potentially modulate PRC2

**Table 2. Statistics for the 20 Energy-Minimized Conformers of PHF1<sub>Tudor</sub> in Complex with H3<sub>31–41</sub>K36Me3**

Distance constraints	
Long ( $ i-j  > 5$ ) (intramolecular)	250
Medium ( $1 <  i-j  \leq 5$ )	106
Sequential ( $ i-j  = 1$ )	234
Intraresidue ( $i = j$ )	462
Intermolecular	38
Dihedral angle constraints ( $\phi$ and $\psi$ )	
	96
Hydrogen bond constraints	
	16
Average pairwise rmsd to the mean structure (Å) <sup>a</sup>	
Backbone (C <sup>α</sup> , C', N, O)	0.26 ± 0.05
Heavy atoms	0.72 ± 0.10
Deviations from idealized covalent geometry	
Bond (Å)	0.008 ± 0.002
Angles (°)	0.744 ± 0.012
Impropers (°)	0.479 ± 0.010
Rmsd from experimental distance restraints (Å)	0.032 ± 0.001
Rmsd from experimental dihedral restraints (°)	0.578 ± 0.027
Ramachandran statistics (% of all residues) <sup>a</sup>	
Most favored	91.3
Additionally allowed	8.7
Generously allowed	0.0
Disallowed	0.0


<sup>a</sup>Statistics are given for residues 31–81 of PHF1<sub>Tudor</sub> and residues 35–40 of the H3<sub>31–41</sub>K36Me3 peptide.

stabilization and spreading during conversion of transcriptionally active/permmissive states to repressed ones—a process often observed in cell differentiation and lineage specification (Mikkelsen et al., 2007). We next tested this hypothesis in HeLa cells, whose endogenous PHF1 expression is low (Boulay et al., 2011), and asked whether ectopic expression of PHF1 could shift chromatin boundaries that separate H3K27me3 from established H3K36me3 domains. Profiling of HeLa cells by chromatin immunoprecipitation coupled with deep sequencing (ChIP-seq) revealed that the *HOX-B* gene clusters, known to be regulated by PRC2 (Cao et al., 2008), exhibited a bimodal distribution of H3K27me3 and H3K36me3 (Figure 4A). H3K27me3 covered the distal *HOX-B* genes (from *HOXB13* to *HOXB7*), and H3K36me3 decorated the proximal ones (from *HOXB6* to *HOXB1*). We first confirmed this bimodal pattern using directed ChIPs (Figures S4A and S4B). In examination of changes in *HOX-B* gene expression before and after ectopic PHF1 expression, we found that WT and Tudor mutant forms of PHF1 demonstrated a chromatin environment-dependent effect on *HOX* repression. First, both WT and Tudor mutant PHF1 repressed *HOXB9*, *HOXB8*, and *HOXB7*, three distal *HOX*s marked by high H3K27me3 and low H3K36me3 (Figures 4A, 4B, and S4C). Using ChIP assays, we detected exogenous FLAG-tagged PHF1 as well as an increase in PRC2 residence and H3K27me3 at *HOXB9* and *HOXB8* (Figures 4C–4E and S4D), suggesting that independent of the Tudor motif, preexisting PRC2 may

recruit PHF1 to H3K27me3-positive regions where PHF1 either enhances PRC2 enzymatic activities as previously reported (Cao et al., 2008; Sarma et al., 2008) or further stabilizes the chromatin binding of PRC2. While we detected no change in expression of proximal *HOX*s positioned deep within the H3K36me3 domain including *HOXB4*, *HOXB3*, and *HOXB1* (Figures 4A and 4B), WT—but not Tudor mutant PHF1—was able to repress *HOXB6* and *HOXB5* (Figure 4B, red line versus green, blue or purple), two *HOX* genes situated exactly on the junction of H3K36me3 and H3K27me3 domains (Figure 4A). These data suggested a Tudor-dependent mechanism for PRC2 targeting to regions of *HOXB6* and *HOXB5*. Indeed, PHF1 residence at *HOXB6* and *HOXB5* was completely dependent on Tudor function (Figure 4C), as was the residence of the PRC2 core complex components EZH2 and SUZ12 and the spread of H3K27me3 into these regions (Figures 4D–4F). No dramatic change in H3K36me3 was detected at examined *HOX-B* loci (Figure S4E). A similar Tudor-dependent H3K27me3 spreading was also observed at *PBX1* and *TADA1/ADA1*, two other developmentally critical genes that also exhibit bimodal distribution of H3K36me3 (at gene bodies) and H3K27me3 (at promoters) (Figures S4F and S4G). Taken together, our data demonstrate that through Tudor-mediated recognition of H3K36me3, ectopically expressed PHF1 is able to mediate the targeting and spreading of associated PRC2 core complexes to H3K36me3-demarcated active chromatin regions to modulate gene expression.

### PHF19/PCL3 Genetic Complementation in Murine Pluripotent Stem Cells Demonstrates a Tudor-Dependent Repression of Differentiation-Associated Genes

We next sought to examine the role of the PCL Tudor motif in PRC2-mediated gene silencing under physiological conditions. Endogenous expression of *PHF1* in HeLa and HEK293 cells is low (Boulay et al., 2011), and its knockdown only induced subtle changes in global H3K27me3 levels and expression of several tested PRC2-regulated genes (data not shown). Since endogenous *PHF1* expression was only found highly expressed in germ cells (Kawakami et al., 1998), we focused on PHF19 (also known as PCL3) (Figure S1B), a murine *PHF1* homolog that is highly expressed in murine ESCs and ESC-like embryonic carcinoma cell line F9 (Figure S5A). Importantly, the Tudor motif of PHF19 showed similar preference and affinity toward H3K36me3 and demonstrated a similar requirement of aromatic caging residues for binding to H3K36me3 peptides (Table 1; Figures S1J, S1K, S2F, and S2G), indicating a functional redundancy among these proteins in recognition of H3K36 methylation. We also found that, unlike a recent study showing that *Phf19* knockdown in ESCs also led to downregulation of *Suz12* (Hunkapiller et al., 2012), *Phf19* knockdown in F9 cells did not cause any detectable change in expression of all examined PRC2 complex genes including *Ezh2*, *Suz12*, and *Mtf2/Pcl2*, as examined by RT-PCR (Figure S5A) and immunoblot (Figure S5B). Knocking down *Phf19* in F9 embryonic carcinoma cells led to derepression of numerous PRC2 direct target genes, as revealed by gene array analysis (Tables S3 and S4) and RT-qPCR (Figure S5A). Consistent with a *Phf19* knockdown



**Figure 3. PHF1<sup>Tudor</sup> Alters Subnuclear Localization of PHF1-PRC2 Complexes, but not PRC2 Core Structure Formation**

(A) Summary of protein peptides identified by mass spectrometry that specifically associate with FLAG-tagged PHF1 complexes purified from 293 stable cell lines. Numbers shown in the table are those of coverage percentage of amino acid sequences, unique peptides, and total peptide hits. See also Table S2.

(B) coIP of FLAG-tagged WT or Tudor-mutated forms of PHF1 with PRC2 core complex components EZH2 and SUZ12. Bottom panels are immunoblots of input. Asterisk represents nonspecific bands. See also Figure S3.

(C) Immunoblot of the indicated FLAG-PHF1 stable expression cell lines separated into chromatin and soluble fractions.


(D) Representative confocal immunofluorescence images revealing localization of WT versus Tudor mutant form of PHF1 proteins (FLAG-tagged, green) and chromatin (stained by DAPI, blue) in transiently transfected 293 cells. Scale bar, 5  $\mu$ m.

experiment performed in E14 murine ESCs (Hunkapiller et al., 2012), the upregulated, but not downregulated, genes found in *Phf19*-knockdown cells were enriched with pathways associated with organismal development and cell differentiation (Tables S3 and S4). Among the top 100 upregulated genes, more than 40% are H3K27me3 positive at their promoters, demonstrating a significant overlap between H3K27me3 and *Phf19*-repressed genes ( $p$  value  $< 2.2 \times 10^{-16}$ , data not shown). Such observed premature derepression of development programs in pluripotent cells is similar to phenotypes observed in *Jarid2* null ESCs (Li et al., 2010; Pasini et al., 2010; Peng et al., 2009; Shen et al., 2009). Thus, F9 embryonic carcinoma cells serve as a suitable model to study functions of the *Phf19* Tudor motif.

Previously, it has been shown that *Phf19* binding sites strongly correlated to those of PRC2 complexes and H3K27me3 peaks (Hunkapiller et al., 2012). Our analysis of ChIP-seq profiling data obtained from murine E14 ESCs (Hunkapiller et al., 2012; Xiao et al., 2012) further revealed a moderate genome-wide association between *Phf19* and H3K36me3, where higher levels of *Phf19* binding were found at promoters of genes with higher levels of H3K36me3 among genes ( $n = 19,387$ ) containing intermediate to low levels of H3K27me3 at promoters (Figure S5C). Analyses of these ESC ChIP-seq data also

revealed a subset of PRC2-targeted genes marked by H3K36me3, which coexisted with H3K27me3 and *Phf19*, as exemplified by *Otx2* (see boxed region, Figure S5D). Using directed ChIP (Figures 5A and S5E–S5H) and sequential re-ChIP analyses (Figure 5B), we confirmed such coexistence of H3K36me3 and H3K27me3 at PRC2 target genes, including *Otx2*, *Meis1*, *Hoxa5*, and *Fgf15*, in both ESCs and F9 cells. At these bivalent PRC-targeted genes, H3K36me3 tends to increase toward gene bodies at the 3' end, and H3K27me3 is enriched at promoter regions in the 5' end. In addition, H3K36me3 was typically detected to a lesser extent when compared to that found at a locus of the housekeeping gene *Rps19* (Figure S5E). Detection of such H3K27me3-H3K36me3 bivalent genes is consistent with recent genomic studies showing coexistence of H3K36me3 and H3K27me3 at multivalent development genes in zebrafish sperm cells (Wu et al., 2011) as well as a proteomic study detecting a small percentage of H3 containing both H3K27 and H3K36 methylation in mammalian cells (Voigt et al., 2012).

Next, we used *Phf19*-expressing F9 cells to examine the function of the H3K36me3-recognizing Tudor motif in regulating the bivalent genes described above. Concurrent with derepression of these development genes following *Phf19* knockdown (Figure S5A), we detected a significant reduction in levels of



**Figure 4. Ectopic Expression of PHF1 Induced a Tudor-Dependent Spreading of PRC2 Complexes and H3K27me3 into an Adjacent H3K36me3 Region within *HOX-B* Gene Clusters**

(A) ChIP-seq revealing a bimodal distribution of H3K27me3 and H3K36me3 at *HOX-B* gene clusters. Position of each *HOX* is indicated at the bottom. See also Figure S4.


(B) Change in expression of *HOX-B* genes in HeLa stable cells with ectopic expression of either WT or Tudor mutant form of PHF1. Overexpression of the WT, but not Tudor mutant form of PHF1, led to repression of *HOXB5* and *HOXB6* (highlighted by a line on the top), two genes situated at the H3K36me3-H3K27me3 conjunction (A). Data of relative mRNA levels (y axis) from three independent experiments were normalized against vector control (shown as a dash line) and then presented as mean  $\pm$  SD.

(C–F) ChIP for FLAG-tagged PHF1 (C), EZH2 (D), SUZ12 (E), and H3K27me3 (F) across different *HOX-B* genes in HeLa cells overexpressing either WT PHF1 or its Tudor mutant form. Data of ChIP signals (y axis) from three independent experiments were normalized to 1% of input used, data of histone modification (F) further normalized to those of total histone H3, and then presented as mean  $\pm$  SD. Statistics shown are t test comparisons of ChIP in cells transduced with WT PHF1 to that with control or the mutant (Y47A) form. \* $p < 0.05$ ; \*\* $p < 0.005$ ; n.s., not significant.

H3K27me3 at all examined target genes (Figures 5C and 5D). No measurable change in H3K36me3 was found at tested loci (Figure S5I). Similar to PHF1 (Figure 3B) and findings of a previous report (Hunkapiller et al., 2012), both WT and Tudor mutant forms of PHF19 efficiently associated with PRC2 complexes (Figure S5J). Introduction of a WT hairpin-resistant PHF19 into F9 cells repressed *Otx2* and other tested target genes (*Meis1*, *Hoxa5*, *Fgf15*) in either parental F9 cells (*Otx2* as tested in Figure 5E) or cells with endogenous Phf19 knocked down (Fig-

ure 5F); the mutants that failed to engage H3K36me3 binding either led to *Otx2* derepression in a dominant-negative fashion in parental F9 cells (Figure 5E) or failed to restore gene repression (Figure 5F). As examined by ChIP assays at all tested gene loci, repression of gene expression was found to be correlated to the binding of WT, but not Tudor mutant PHF19 (Figure 5G), the recruitment of PRC2 core components (Figure 5H), and the level of H3K27me3 (Figure S5K). Collectively, our data suggest that PHF19, another member of PCL family





**Figure 5. Phf19/Pcl3 Demonstrates a Tudor-Dependent Repression of Differentiation-Associated Genes**

(A) ChIP for H3K36me3 and H3K27me3 across the *Otx2* gene in F9 embryonic carcinoma cells. The diagram on top depicts the genomic organization of *Otx2* with positions of each ChIP PCR amplicon marked as (a–h) (not drawn to scale). Data of ChIP signals (y axis) from three independent experiments were normalized to 1% of input and then to histone H3 and presented as mean  $\pm$  SD. See also Figure S5.

(B) Sequential re-ChIP showing fold of enrichment by comparing ChIP signals of H3K27me3 over IgG control in a sequential IP following the first IP using H3K36me3 antibodies. Signal of ChIP (y axis) from three independent experiments were normalized to 1% of input and then to nonspecific IgG control and presented as mean  $\pm$  SD. Statistics shown are t test comparisons of each tested loci to three negative control loci, the transcriptional start site (TSSs) of *Oct4* and  $\beta$ -actin, and the gene body of *Rps19*. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.005$ ; n.s., not significant.

(C and D) ChIP examining H3K27me3 levels associated with *Otx2* (C) and other examined development genes (D) in F9 cells transduced with control or *Phf19*-specific hairpins. Data of ChIP signals (y axis) from three independent experiments were normalized to 1% of input and then to histone H3 and presented as mean  $\pm$  SD. ex, exon; TES, transcriptional ending site.

(E) Real-time PCR detecting *Otx2* expression levels in F9 cells following overexpression of WT and Tudor mutant forms of *PHF19/PCL3*. Data of relative mRNA levels (y axis) from three independent experiments were normalized to vector transduced cells and presented as mean  $\pm$  SD.

(F) Real-time PCR of *Otx2*, *Meis1*, *Hoxa5*, *Fgf15*, and *Oct4* expression in F9 cells following *Phf19* knockdown (sh\_*Phf19*) and reintroduction of hairpin-resistant WT and Tudor mutant forms of FLAG-tagged human PHF19/PCL3 (F-PHF19). y axis represents fold change after normalization of data from three independent experiments to control cells (presented as mean  $\pm$  SD). See also Tables S3 and S4.

(G and H) ChIP examining the residence of FLAG-tagged PHF19/PCL3 (G) and SUZ12 (H) at *Otx2*, *Meis1*, *Hoxa5*, and *Fgf15* genes after transduction of hairpin-resistant WT or Tudor mutant PHF19/PCL3 into knockdown cells. Signal of ChIP (y axis) from three independent experiments were normalized to 1% of input and presented as mean  $\pm$  SD. IgG and TSS of  $\beta$ -actin were used as antibody and locus control, respectively. Statistics shown are t test comparisons of ChIP in cells transduced with WT PHF19 to that with Tudor mutant forms. \* $p < 0.01$ ; \*\* $p < 0.001$ ; \*\*\* $p < 0.0001$ .

proteins, also regulates repression of PRC2 targets via a functional Tudor motif that recognizes H3K36me3.

## DISCUSSION

### PCL Family Identified as High-Affinity H3K36me3 Effector Proteins

In this study, we have identified the PCL family of Tudor motifs as H3K36me3 effector modules. Our structural, biochemical, and biophysical studies of these Tudor motifs revealed an aromatic H3K36me3-caging pocket that forms direct contacts to H3K36me3 as well as to the surrounding H3 residues, which explained our observed high affinity and selectivity. In comparison to the previously identified H3K36me3-reading effectors BRPF1 and EAF3, whose affinities to H3K36me3 were measured in the high  $\mu\text{M}$  to low mM range ( $K_D = \sim 0.3\text{--}3\text{ mM}$ ), both the PHF1 and PHF19 Tudor motifs bind to H3K36me3 with affinities in the low  $\mu\text{M}$  range ( $K_D = \sim 5\text{--}7\ \mu\text{M}$ ). These  $K_D$  measurements are similar to affinities measured for effectors of other common histone methylation marks, including H3K4me3-recognizing PHD fingers ( $K_D = \sim 1\text{--}10\ \mu\text{M}$ ) (Li et al., 2006; Wang et al., 2009), H3K27me3- and H3K9me3-recognizing chromo domains ( $K_D = \sim 5\text{--}50\ \mu\text{M}$ ) (Bernstein et al., 2006; Kaustov et al., 2011), and an H3K9me3-recognizing Tudor domain (Rothbart et al., 2012a). Notably, PCL family proteins also contain two PHD fingers, one of which is in close proximity to the Tudor motif (Figure S1B); the histone-binding activities of these domains, as well as their influence on PCL functions, remain to be determined.

### PCL Proteins Serve as a Bridging and/or Targeting Mechanism for PRC2 to Intrude into Active Chromatin Regions

Our collective biochemical and genetic studies demonstrate a PRC2-targeting mechanism wherein the cofactor PCL proteins either stabilize, tether, or establish intrusion of PRC2 core components into the H3K36me3-containing chromatin via Tudor-mediated recognition of H3K36me3 and histone H3. coIP studies of PHF1 showed that this protein associates with PRC2 core components and assembles a higher-order complex efficiently, a process independent of the Tudor motif. We further show that PCL proteins promote PRC2 chromatin association and catalysis of H3K27me3, consistent with previous observations (Cao et al., 2008; Hunkapiller et al., 2012; Sarma et al., 2008). Our studies of ectopic PHF1 expression, modulation of global H3K36me3, as well as knockdown of endogenous Phf19 (without changing PRC2 core complexes per se) demonstrated a role of PCL proteins in the bridging of PRC2 to H3K36me3-containing nucleosomes, a mechanism that may act together with those provided by other PRC2-targeting mechanisms such as EED (Margueron et al., 2009), JARID2 (Li et al., 2010; Pasini et al., 2010; Peng et al., 2009; Shen et al., 2009), and noncoding RNAs (Bracken and Helin, 2009; Margueron and Reinberg, 2011). As we did not detect association of PHF1 or PHF19 to non-PRC2 target genes that are highly H3K36me3 positive (exemplified by *CD44* in Figure S4H and  $\beta$ -actin in Figure 5G), an intriguing model would be that noncoding RNA, DNA-binding, and chromatin-binding activities associated with

PRC2 complexes and cofactors (for example EED and JARID2) collectively mediate initial PRC2 recruitment to H3K27me3-positive promoters/nucleosomes, while PCL proteins further modulate PRC2 stabilization and tether PRC2 to adjacent H3K36me3/H3K36me2-positive nucleosomes toward gene bodies. Although it has been previously shown that active histone methylation marks, including H3K36me3, inhibit PRC2-mediated H3K27me3 (Schmitges et al., 2011; Yuan et al., 2011), our studies demonstrate that PRC2 spreads and intrudes into active chromatin during gene silencing and chromatin remodeling, a critical process concurrent with development and differentiation (Mikkelsen et al., 2007).

### Coexisting Together: How Do Seemingly Biologically Opposing Chromatin Marks Work Together in the Chromatin Landscape?

Our genomics and ChIP analysis revealed that relatively lower levels of H3K36me3 exist in previously described poised bi- or multivalent chromatin state genes found in ESCs (Mikkelsen et al., 2007; Xiao et al., 2012). Such detected H3K36me3 may represent residual species of H3K36me3 docked by Phf19 in processes of PRC2-mediated chromatin remodeling associated with lineage differentiation and development. Indeed, a recent mass spectrometry-based analysis identified asymmetrically methylated, bivalent mononucleosomes that carry H3K36me3/H3K36me2 along with H3K27me3 (Voigt et al., 2012). We suspect that these asymmetrically methylated mononucleosomes are more likely to be found at conjunction regions of H3K27me3 and H3K36me3 in those bivalent or bimodal genes or loci as examined in our study (exemplified by *HOX-B* gene cluster in Figure 4A, *PBX1* in Figure S4F, and *Otx2* in Figure 5A) where PCL proteins serve as a bridging and stabilizing mechanism for PRC2 complex encroachment and spreading.

We also observed overall separation of high peaks/domains of H3K27me3 and H3K36me3 in ChIP-seq as previously reported (Mikkelsen et al., 2007; Xiao et al., 2012) (data not shown), and we suspect that PCL proteins may be involved in a dynamic process aimed at conversion of active chromatin state to the poised/repressed state by recruiting additional factors such as H3K36 demethylases to achieve completion of remodeling. We also hypothesize that residual H3K36me3 marks retained at these poised development genes might serve as a memory of cellular and epigenetic states, a mechanism that separates them from those silenced permanently. In accordance, expression of PCL family genes is dynamically regulated during organismal development, with *PHF1* specifically expressed in murine germ cells (Kawakami et al., 1998) and *Mtf2/Pcl2* preferentially expressed in undifferentiated ESCs (Walker et al., 2010). Both *PHF1* and *PHF19* were reported to be translocated or overexpressed in various human tumor types (Micci et al., 2006; Wang et al., 2004). Collectively, our studies strongly indicate that the chromatin- and H3K36me3-recognizing activities harbored in mammalian PCL proteins are critical for PRC2 targeting and regulation of developmentally critical genes. They further shed light on the modulation of the chromatin landscape and gene expression dynamics during processes of development and oncogenesis.

## EXPERIMENTAL PROCEDURES

### Plasmid Construction

Plasmids containing cDNA of PHF1 (also known as polycomb-like 1 or PCL1, NCBI accession NM\_024165) and PHF19 (also known as polycomb-like 3 or PCL3, NCBI accession BC125076) were purchased from Open Biosystems. cDNAs were fused to either HA-FLAG or 2xFLAG tag by PCR and cloned into MSCV retroviral expression vectors (Clontech) and pGEX GST-fusion plasmids (GE Life Science). Deletion and mutation were introduced by PCR and site-directed mutagenesis. All plasmids used were confirmed by sequencing.

### Recombinant Glutathione S-Transferase Protein Production

Glutathione S-transferase (GST)-fusion proteins were generated and purified as previously described (Ruthenburg et al., 2011; Wang et al., 2009).

### Histone Peptide Microarrays

Methods for fabrication of the histone peptide array platform (with peptide identities listed in Table S1) and analysis of effector protein binding were previously described (Fuchs et al., 2011; Rothbart et al., 2012b). A second histone peptide array platform was also obtained and used according to the manufacturer's specifications (Active Motif).

### Chromatin Fractionation

Chromatin and associated proteins were isolated from asynchronously growing 293 cells as described (Rothbart et al., 2012a).

### ChIP and Sequential Re-ChIP Assays

ChIP was performed using a previously described protocol (Wang et al., 2009). ChIP signals were represented as a percentage of input chromatin, and fold of enrichment was calculated by normalizing against signals of nonspecific IgG. Sequential re-ChIP was performed using a previously described protocol (Voigt et al., 2012). Information regarding primers used is described in Table S5.

### NMR Spectroscopy

Collection of NMR spectra and structural calculations are detailed in Supplemental Experimental Procedures.

## ACCESSION NUMBERS

The structural coordinates of PHF1 Tudor domain protein in the complex with H3<sub>31-41</sub>K36me3 peptide have been deposited in the Protein Data Base under accession number 2M0O. The microarray data of Phf19 shRNA knockdown have been deposited in GEO database under accession number GSE42463.

## SUPPLEMENTAL INFORMATION

Supplemental Information includes five figures, five tables, and Supplemental Experimental Procedures and can be found with this article online at <http://dx.doi.org/10.1016/j.molcel.2012.11.026>.

## ACKNOWLEDGMENTS

We thank K. Krajewski for synthesizing peptides, M. Vernon for gene microarray analysis, Y. Zhang for providing JMJD2 plasmids, and T. Magnuson for anti-EED antibodies used in this study. L.C. and S.B.R. are supported by postdoctoral fellowships awarded by the US Army-DOD Prostate Cancer Research Program and the UNC Lineberger Comprehensive Cancer Center Basic Sciences Training Program (T32CA09156), respectively. This research was also supported by grants from the Leukemia & Lymphoma Society and STARR Foundation to both D.J.P. and C.D.A., and National Institutes of Health (NIH) grants to B.D.S. (GM068088). J.S. is supported by funds from UC-Riverside, and G.G.W. is supported by the NIH/NCI "Pathway to Independence" Award in Cancer Research (CA151683) and University Cancer Research Fund (UCRF)

of the NC State. G.G.W. is a Martin D. Abeloff, M.D. V Scholar of the V Foundation for Cancer Research.

Received: September 20, 2012

Revised: October 31, 2012

Accepted: November 21, 2012

Published: December 27, 2012

## REFERENCES

- Bernstein, E., Duncan, E.M., Masui, O., Gil, J., Heard, E., and Allis, C.D. (2006). Mouse polycomb proteins bind differentially to methylated histone H3 and RNA and are enriched in facultative heterochromatin. *Mol. Cell Biol.* 26, 2560–2569.
- Boulay, G., Rosnoblet, C., Guérardel, C., Angrand, P.O., and Leprince, D. (2011). Functional characterization of human Polycomb-like 3 isoforms identifies them as components of distinct EZH2 protein complexes. *Biochem. J.* 434, 333–342.
- Bracken, A.P., and Helin, K. (2009). Polycomb group proteins: navigators of lineage pathways led astray in cancer. *Nat. Rev. Cancer* 9, 773–784.
- Cao, R., and Zhang, Y. (2004). SUZ12 is required for both the histone methyltransferase activity and the silencing function of the EED-EZH2 complex. *Mol. Cell* 15, 57–67.
- Cao, R., Wang, H., He, J., Erdjument-Bromage, H., Tempst, P., and Zhang, Y. (2008). Role of hPHF1 in H3K27 methylation and Hox gene silencing. *Mol. Cell Biol.* 28, 1862–1872.
- Chi, P., Allis, C.D., and Wang, G.G. (2010). Covalent histone modifications—miswritten, misinterpreted and mis-erased in human cancers. *Nat. Rev. Cancer* 10, 457–469.
- Friberg, A., Oddone, A., Klymenko, T., Müller, J., and Sattler, M. (2010). Structure of an atypical Tudor domain in the Drosophila Polycomblike protein. *Protein Sci.* 19, 1906–1916.
- Fuchs, S.M., Krajewski, K., Baker, R.W., Miller, V.L., and Strahl, B.D. (2011). Influence of combinatorial histone modifications on antibody and effector protein recognition. *Curr. Biol.* 21, 53–58.
- Hunkapiller, J., Shen, Y., Diaz, A., Cagney, G., McCleary, D., Ramalho-Santos, M., Krogan, N., Ren, B., Song, J.S., and Reiter, J.F. (2012). Polycomb-like 3 promotes polycomb repressive complex 2 binding to CpG islands and embryonic stem cell self-renewal. *PLoS Genet.* 8, e1002576.
- Kaustov, L., Ouyang, H., Amaya, M., Lemak, A., Nady, N., Duan, S., Wasney, G.A., Li, Z., Vedadi, M., Schapira, M., et al. (2011). Recognition and specificity determinants of the human cbx chromodomains. *J. Biol. Chem.* 286, 521–529.
- Kawakami, S., Mitsunaga, K., Kikuti, Y.Y., Ando, A., Inoko, H., Yamamura, K., and Abe, K. (1998). Tctex3, related to Drosophila polycomblike, is expressed in male germ cells and mapped to the mouse t-complex. *Mamm. Genome* 9, 874–880.
- Klose, R.J., and Zhang, Y. (2007). Regulation of histone methylation by demethylination and demethylation. *Nat. Rev. Mol. Cell Biol.* 8, 307–318.
- Kouzarides, T. (2007). Chromatin modifications and their function. *Cell* 128, 693–705.
- Li, H., Ilin, S., Wang, W., Duncan, E.M., Wysocka, J., Allis, C.D., and Patel, D.J. (2006). Molecular basis for site-specific read-out of histone H3K4me3 by the BPTF PHD finger of NURF. *Nature* 442, 91–95.
- Li, G., Margueron, R., Ku, M., Chambon, P., Bernstein, B.E., and Reinberg, D. (2010). Jarid2 and PRC2, partners in regulating gene expression. *Genes Dev.* 24, 368–380.
- Margueron, R., and Reinberg, D. (2011). The Polycomb complex PRC2 and its mark in life. *Nature* 469, 343–349.
- Margueron, R., Justin, N., Ohno, K., Sharpe, M.L., Son, J., Drury, W.J., 3rd, Voigt, P., Martin, S.R., Taylor, W.R., De Marco, V., et al. (2009). Role of the polycomb protein EED in the propagation of repressive histone marks. *Nature* 461, 762–767.

- Micci, F., Panagopoulos, I., Bjerkehagen, B., and Heim, S. (2006). Consistent rearrangement of chromosomal band 6p21 with generation of fusion genes JAZF1/PHF1 and EPC1/PHF1 in endometrial stromal sarcoma. *Cancer Res.* *66*, 107–112.
- Mikkelsen, T.S., Ku, M., Jaffe, D.B., Issac, B., Lieberman, E., Giannoukos, G., Alvarez, P., Brockman, W., Kim, T.K., Koche, R.P., et al. (2007). Genome-wide maps of chromatin state in pluripotent and lineage-committed cells. *Nature* *448*, 553–560.
- Pasini, D., Cloos, P.A., Walfridsson, J., Olsson, L., Bukowski, J.P., Johansen, J.V., Bak, M., Tommerup, N., Rappsilber, J., and Helin, K. (2010). JARID2 regulates binding of the Polycomb repressive complex 2 to target genes in ES cells. *Nature* *464*, 306–310.
- Peng, J.C., Valouev, A., Swigut, T., Zhang, J., Zhao, Y., Sidow, A., and Wysocka, J. (2009). Jarid2/Jumonji coordinates control of PRC2 enzymatic activity and target gene occupancy in pluripotent cells. *Cell* *139*, 1290–1302.
- Rothbart, S.B., Krajewski, K., Nady, N., Tempel, W., Xue, S., Badeaux, A.I., Barsyte-Lovejoy, D., Martinez, J.Y., Bedford, M.T., Fuchs, S.M., et al. (2012a). Association of UHRF1 with methylated H3K9 directs the maintenance of DNA methylation. *Nat. Struct. Mol. Biol.* *19*, 1155–1160.
- Rothbart, S.B., Krajewski, K., Strahl, B.D., and Fuchs, S.M. (2012b). Peptide microarrays to interrogate the “histone code”. *Methods Enzymol.* *512*, 107–135.
- Ruthenburg, A.J., Li, H., Milne, T.A., Dewell, S., McGinty, R.K., Yuen, M., Ueberheide, B., Dou, Y., Muir, T.W., Patel, D.J., and Allis, C.D. (2011). Recognition of a mononucleosomal histone modification pattern by BPTF via multivalent interactions. *Cell* *145*, 692–706.
- Sarma, K., Margueron, R., Ivanov, A., Pirrotta, V., and Reinberg, D. (2008). Ezh2 requires PHF1 to efficiently catalyze H3 lysine 27 trimethylation in vivo. *Mol. Cell Biol.* *28*, 2718–2731.
- Schmitges, F.W., Prusty, A.B., Faty, M., Stützer, A., Lingaraju, G.M., Aiwazian, J., Sack, R., Hess, D., Li, L., Zhou, S., et al. (2011). Histone methylation by PRC2 is inhibited by active chromatin marks. *Mol. Cell* *42*, 330–341.
- Selenko, P., Sprangers, R., Stier, G., Bühler, D., Fischer, U., and Sattler, M. (2001). SMN tudor domain structure and its interaction with the Sm proteins. *Nat. Struct. Biol.* *8*, 27–31.
- Shen, X., Kim, W., Fujiwara, Y., Simon, M.D., Liu, Y., Mysliwiec, M.R., Yuan, G.C., Lee, Y., and Orkin, S.H. (2009). Jumonji modulates polycomb activity and self-renewal versus differentiation of stem cells. *Cell* *139*, 1303–1314.
- Sun, B., Hong, J., Zhang, P., Dong, X., Shen, X., Lin, D., and Ding, J. (2008). Molecular basis of the interaction of *Saccharomyces cerevisiae* Eaf3 chromo domain with methylated H3K36. *J. Biol. Chem.* *283*, 36504–36512.
- Vermeulen, M., Eberl, H.C., Matarese, F., Marks, H., Denissov, S., Butter, F., Lee, K.K., Olsen, J.V., Hyman, A.A., Stunnenberg, H.G., and Mann, M. (2010). Quantitative interaction proteomics and genome-wide profiling of epigenetic histone marks and their readers. *Cell* *142*, 967–980.
- Vezzoli, A., Bonadies, N., Allen, M.D., Freund, S.M., Santiveri, C.M., Kvinlaug, B.T., Huntly, B.J., Göttgens, B., and Bycroft, M. (2010). Molecular basis of histone H3K36me3 recognition by the PWWP domain of Brpf1. *Nat. Struct. Mol. Biol.* *17*, 617–619.
- Voigt, P., LeRoy, G., Drury, W.J., 3rd, Zee, B.M., Son, J., Beck, D.B., Young, N.L., Garcia, B.A., and Reinberg, D. (2012). Asymmetrically modified nucleosomes. *Cell* *151*, 181–193.
- Walker, E., Chang, W.Y., Hunkapiller, J., Cagney, G., Garcha, K., Torchia, J., Krogan, N.J., Reiter, J.F., and Stanford, W.L. (2010). Polycomb-like 2 associates with PRC2 and regulates transcriptional networks during mouse embryonic stem cell self-renewal and differentiation. *Cell Stem Cell* *6*, 153–166.
- Wang, S., Robertson, G.P., and Zhu, J. (2004). A novel human homologue of *Drosophila* polycomblike gene is up-regulated in multiple cancers. *Gene* *343*, 69–78.
- Wang, G.G., Song, J., Wang, Z., Dormann, H.L., Casadio, F., Li, H., Luo, J.L., Patel, D.J., and Allis, C.D. (2009). Haematopoietic malignancies caused by dysregulation of a chromatin-binding PHD finger. *Nature* *459*, 847–851.
- Wu, S.F., Zhang, H., and Cairns, B.R. (2011). Genes for embryo development are packaged in blocks of multivalent chromatin in zebrafish sperm. *Genome Res.* *21*, 578–589.
- Xiao, S., Xie, D., Cao, X., Yu, P., Xing, X., Chen, C.C., Musselman, M., Xie, M., West, F.D., Lewin, H.A., et al. (2012). Comparative epigenomic annotation of regulatory DNA. *Cell* *149*, 1381–1392.
- Xu, C., Cui, G., Botuyan, M.V., and Mer, G. (2008). Structural basis for the recognition of methylated histone H3K36 by the Eaf3 subunit of histone deacetylase complex Rpd3S. *Structure* *16*, 1740–1750.
- Yuan, W., Xu, M., Huang, C., Liu, N., Chen, S., and Zhu, B. (2011). H3K36 methylation antagonizes PRC2-mediated H3K27 methylation. *J. Biol. Chem.* *286*, 7983–7989.